



Received for publication, June, 22, 2017

Accepted, January, 5, 2018

## *Original paper*

# **Research on evolution of nitrite and nitrate content regarding milk processing in scalded cheese**

DORIN ȚIBULCĂ<sup>1</sup>, MIRELA JIMBOREAN<sup>\*1</sup>, AURORA ȚIBULCĂ<sup>2</sup>

<sup>1</sup>University of Agricultural Sciences and Veterinary Medicine, Faculty of Food Science and Technology, Department of Food Engineering, 3-5 Mănăștur Street, Cluj-Napoca, România

<sup>2</sup>S.C. RAAL S.A. Bistrița, România

### **Abstract**

Nitrates have an insignificant toxicity. They are considered virtual toxic substance because they convert into nitrogen, which have a toxic potential well known. Milk contaminated with nitrates must be severely restricted.

Objectives of our research took into account the wholesomeness of milk to determine nitrates and nitrites. Sixty samples of raw milk (15 samples per season) were taken into consideration, to see the level of nitrites and nitrates from processed milk for consumption. To study the remanence of nitrates and nitrites in the finished product 60 samples of cheese were also analyzed.

Following the conducted research, no evidence has exceeded the maximum permissible limit for nitrates and nitrites which is max. 10 mg/ liter milk. For samples of cheese were not found values above 10 ppm that is the maximum permissible limit for nitrates and nitrites in cheese. Nitrates are concentrated in cheese 1.3 times. The content of nitrites in the cheese falls by 1.9 times, from 0.03367 mg/ l in milk to 0.01728 mg/ kg in cheese.

### **Keywords**

Residues of nitrites and nitrates, Griess method, colorimetric method with m-xyleneol, cheese.

**To cite this article:** ȚIBULCĂ D, JIMBOREAN M, ȚIBULCĂ A. Research on evolution of nitrite and nitrate content regarding milk processing in scalded cheese. *Rom Biotechnol Lett.* 2019; 24(5): 770-775. DOI: 10.25083/rbl/24.5/770.775

---

✉ \*Corresponding author: MIRELA JIMBOREAN, University of Agricultural Sciences and Veterinary Medicine, Faculty of Food Science and Technology, Department of Food Engineering, 3-5 Mănăștur Street, Cluj-Napoca, România  
E-mail: [mirela.jimborean@usamvcluj.ro](mailto:mirela.jimborean@usamvcluj.ro)

## Introduction

Nitrates and nitrites added to food can cause formation of cancerous N-nitroso compounds, whereas exposure to perchlorate is especially emphasised as an important risk factor for newborns' health. (SUNGUR [17]) Some epidemiological studies linking intake of nitrate and nitrite with gastric cancer in humans indicated a positive correlation. (SIMION & al [16]).

The main sources of contamination of milk are: water, placing fraudulent ammonium nitrate in milk and nitric acid used to remove "Stone Milk" on pasteurizers with plate. Nitrites from milk are highly toxic, especially for children. Nitrites concentrations grow 5-10 times in severe heat treatment, therefore, to feed infants and children separation techniques through semi-permeable membrane were developed. (ȚIBULCĂ [21]).

Water polluted with nitrates and nitrites can get into milk especially during cleaning operations of equipment and containers. The main source of water contamination is the ammonium nitrate used as basic chemical fertilizer in agricultural practice (SINDILAR [18], JIMBOREAN & al [11]).

Nitrates and nitrites get in animals through feed and water. Nitrogen is absorbed by plants in the form of either ammonium ( $\text{NH}_4^+$ ) or nitrate ( $\text{NO}_3^-$ ), and its accumulation is influenced by a series of factors that are depending on the species, cultivar, age and soil conditions (SIMION & al [16]). Although animals can ingest large amounts of nitrates and nitrites, organic tissues contain relatively small amounts, due to rapid metabolism and excretion in the urine. Additionally, in ruminant a large part of nitrates and nitrites is subject to changes by ruminal microflora. (BONDOC [2], COSTIN [7]). The nitrate may naturally present in milk and the level of it depends on the quality of feeding materials (i.e. water, feeds) of cows. (TOPÇU [19]).

Nitrogen content in milk is very low. According to Walker, milk content in nitrogen is between the values of 20 and 40 mg nitrate/ liter. This variation in the analytical data is determined by nature of the feed consumed. (WALKER [23]).

In preparations of milk in 2006, the Brasov area, the average content of nitrite was 0.009 g/ kg and ranged from not detected to 0.64 mg/ kg, and the average nitrate content was 0.95 mg/ kg and ranged from not detected and 12.95 mg/ kg. (CRISTEA [4]).

In some countries may be added nitrate in the milk used to prepare cheese to prevent bloating caused by coliform, but mainly to prevent latter bloating produced by some clostridia. Part of the nitrate is removed in the whey and some are converted into harmless compounds in the maturation process. (COSTIN [6], JIMBOREAN [10]) In addition, the residual quantity decreases gradually during maturation (GUŞ [8], BONDOC [2]).

In Brazil, the use of these additives is permitted at maximum levels of 50 mg/kg. The basis of the previously validated method is the quantitative reduction of nitrate to

nitrite through cadmium column and spectrophotometric determination after nitrite diazotation with sulphanilic acid/alpha-naphthol reagent. From all samples analyzed, 38 (88.37%) showed neither a nitrate nor nitrite detectable content; 5 samples (11.63%) presented nitrate, 4 of them being above the tolerated level and only one (2.33%) showed detectable nitrite, however below the permissible limit. The majority of samples did not show nitrate or nitrite at detectable levels. However, though probably safe from a toxicological point of view, the results can implicate an increased risk of bacteriological contamination. (SERAPHIM [15]). In other researches, the highest level of nitrate and nitrite was found in green cheese (17.52 mg/kg and respectively at 21.16 mg/kg) obtained from sheep milk. (TUDOR & al [20]).

Objectives of our research are to assess the level of nitrites and nitrates from processed milk for their consumption and remains detectable in the final product: cheese.

## Materials and Methods

### Cheese preparation

Cheeses have the primary feature of the technological process to obtain curd which, after maturation and slicing, is subjected to scalding temperature of 72-74°C (BANU [1] CODOBAN [5], COSTIN [6], ȚIBULCĂ [22], JIMBOREAN [10]).

Curd preparation involves the following: milk reception, cleaning and standardization of milk, milk pasteurization and cooling, preparation for coagulation which consists of adding calcium chloride (clotting ability of milk increases, it adds 10 to 20 g/100 liters of milk) and selected cultures of lactic bacteria (*Lactococcus lactis* ssp. *lactis*, *Streptococcus thermophilus* and *Lactobacillus casei*). *Lactococcus lactis* ssp. *lactis* is a regular agent of milk spontaneous acidification, hydrolyzing lactose into lactic acid. *Streptococcus thermophilus* produce a weak acidification and a specific flavor. *Lactobacillus casei* is a proteolytic bacteria and plays a role in cheese maturation. (BANU [1], CODOBAN [5], COSTIN [6], ȚIBULCĂ [22], JIMBOREAN [10]). Pasteurization is able to destroy essential microflora, enzymes and pathogens in milk. It should be noted that inactivation level of microorganisms depends on the amount of microorganisms, growth phase and other factors. (CIPROVICA [3]).

The coagulation of milk takes place at 32 ... 35°C, for 30-40 minutes. Processing the curd is then made, and the resulting curd matures at 18-26°C for 6 ... 10 hours after a pH 4.8 to 5.0 is achieved in curd. Mature curd is cut into slices having a thickness of 0.3 to 0.5 cm and a width of 3-6 cm and is scalded at 72 ... 74°C for 50-60 s in brine with 10-12% salt. The slurry is allowed to air in molds, and is matured. Storage is done at 4 ... 6°C and  $\phi = 85-90\%$ . (GUZUN [9]).

### Experimental design

The reference method for the determination of nitrite in milk is Griess method. The reference method for nitrates is colorimetric method with meta-xylenol. This method determines total nitrogen. Nitrate content is calculated by the difference between total nitrogen and the nitrite determined by Griess method and expressed in equivalent nitrate. (JIMBOREAN & al [12], MUREȘAN [14]).

Determination of nitrate and nitrite is done through the following methods: determination of nitrate with diphenylamine (nitrates form with diphenylamine a blue colored complex; color intensity is proportional to the concentration of nitrates in the analyzed sample and is measured colorimetrically); the method for determining the indole nitrite (nitrite formed with indole in sulfuric acid medium, a pink-colored complex; the intensity of the color is proportional to the nitrite concentration in the sample to be analyzed and is measured colorimetrically). In case of dispute the method used is to reduce nitrate to nitrite in cadmium copper environment. The reagents used must be of quality for analysis or of equivalent quality. Water must be distilled or of equivalent purity, free of nitrates and nitrites. (ȚIBULCĂ [21]).

### Results and Discussions

$\text{NO}_3^-$  and  $\text{NO}_2^-$  limits for milk and measures to be taken are:

For  $\text{NO}_3^-$ :

- lack  $\text{NO}_3^-$ , but correlated with  $\text{NO}_2^-$  it is freely admitted in consumption;
- up to 10 mg/ liter is allowed in consumption without restrictions;
- from 10 mg/ liter to 100 mg/ liter is excluded from food for children under one year and is admitted for adults;
- more than 100 mg/ liter is excluded from consumption and is processed in fermented cheese products.

For  $\text{NO}_2^-$ :

- lack  $\text{NO}_2^-$  but correlated with  $\text{NO}_3^-$  is freely allowed freely in consumption;
- less than 10 mg/ liter is allowed without restrictions in consumption;
- more than 10 mg/ liter, regardless of the amount of  $\text{NO}_3^-$  is excluded from consumption and is processed in fermented chesses. (BONDOC [2])

The determined levels of nitrates and nitrites in raw milk are shown in Table 1.

**Table 1.** Variation of nitrates and nitrites content in raw milk

Crt. No	Season	Samples/season	Nitrates $\text{NO}_3^-$ , mg/l			Nitrites $\text{NO}_2^-$ , mg/l		
			0-0.05	0.051-0.1	> 0.1	0-0.05	0.051-0.1	> 0.1
1.	A	15	3	12	-	10	5	-
2.	W	15	-	14	1	8	7	-
3.	Sp	15	4	11	-	13	2	-
4.	Su	15	1	14	-	13	2	-
TOTAL		60	8	51	1	44	16	-
%		100	13.3	85	1,7	73.3	26.7	-

legend: A/ autumn; W/ winter; Sp/ spring; Su/ summer.

Of the 60 milk samples investigated no evidence has exceeded the maximum permissible limit for nitrates and nitrites which is max. 10 mg/l milk. Nitrates had an average ( $\pm$ ) of  $0.0623 \pm 0.004678$  mg/l and a coefficient of variation of 58.16%. The determined values ranged between 0.001 and 0.3 mg/l. Statistically analyzing the nitrates variation, per seasons, shows the following:

- In autumn, nitrates, had an average ( $\pm$ ) of  $0.05287 \pm 0.006306$  mg/l and a coefficient of variation of 46.19%. The determined values ranged between 0.001 and 0.092 mg/l.

- In winter, nitrates, had an average ( $\pm$ ) of  $0.08867 \pm 0.01547$  mg/l and a coefficient of variation of 67.57%. The determined values ranged between 0.057 and 0.3 mg/l.

- In spring, nitrates, had an average ( $\pm$ ) of  $0.0552 \pm 0.004762$  mg/l and a coefficient of variation of

33.41%. The determined values ranged between 0.001 and 0.092 mg/l.

- In summer, nitrates, had an average ( $\pm$ ) of  $0.0007488 \pm 0.05247$  mg/l and a coefficient of variation of 5.53%. The determined values ranged from 0.05 to 0.061 mg/l.

Nitrites had an average ( $\pm$ ) of  $0.03367 \pm 0.003034$  mg/l and a coefficient of variation of 69.81%. The determined values ranged between 0.0 and 0.08 mg/l. Statistically analyzing the nitrites variation, per seasons, shows the following:

- In autumn, nitrites, had an average ( $\pm$ )  $0.036 \pm 0.007024$  mg/l and a coefficient of variation of 75.56%. The determined values ranged between 0.01 and 0.08 mg/l.

- In winter, nitrites, had an average ( $\pm$ ) of  $0.05 \pm 0.004781$  mg/l and a coefficient of variation of 37.03%. The determined values ranged between 0.02 and 0.08 mg/l.

- In spring, nitrites, had an average ( $\pm$ ) of  $0.02667 \pm 0.00504$  mg/l and a coefficient of variation of 73.19%. The determined values ranged between 0.01 and 0.07 mg/l.

- In summer, nitrites, had an average ( $\pm$ )  $0.022 \pm 0.004995$  mg/l and a coefficient of variation of 87.94%. The determined values ranged between 0.0 and 0.06 mg/l.

In Romania it is forbidden the use of nitrite in milk and milk products. Nitrites and nitrates are not toxic in their normal concentrations in food. Exceeding certain limits, however, have serious repercussions on the human body (COSTIN [7]).

A large study whose results were presented by Şindilar (ŞINDILAR [18]) included a total of 210 milk samples, of

which 95 samples of collected milk, where nitrates had limits of variation between 0 and 7.5 mg  $\text{NO}_3^-$ /l, with an average of 2.7  $\text{NO}_3^-$ /l.

The limits of variation of nitrate and nitrite in studied cheese samples are shown in Tables 2 and 3.

As shown in table 2 values of nitrate were recorded as follows:

- 93.3% of cheese samples had values of nitrate between 0-0.1 ppm;
- 6.7% of cheese samples had values of nitrate between 0.11 to 1 ppm.

**Table 2.** Variation limits of nitrates content in pressed cheese samples investigated

Crt. No	Season	Samples	Nitrates, ppm							
			0-0.1	%	0.11-1	%	1.1-10	%	> 10	%
1.	A	15	14	93.3	1	6.7	-	-	-	-
2.	W	15	9	60	6	40	-	-	-	-
3.	Sp	15	14	93.3	1	6.7	-	-	-	-
4.	Su	15	15	100	-	-	-	-	-	-
Total		60	52	86.7	8	13.3	-	-	-	-

legend: A/ autumn; W/ winter; Sp /spring; Su/ summer

**Table 3.** Variation limits of nitrites content in pressed cheese samples investigated

Crt. No	Season	Samples	Nitrites, ppm							
			0-0.01	%	0.011-0.1	%	0.11-10	%	> 10	%
1.	A	15	8	53.3	7	46.7	-	-	-	-
2.	W	15	4	26.7	11	73.3	-	-	-	-
3.	Sp	15	8	53.3	7	46.7	-	-	-	-
4.	Su	15	8	53.3	7	46.7	-	-	-	-
Total		60	28	46.7	32	53.3	-	-	-	-

legend: A/ autumn; W/ winter; Sp /spring; Su/ summer

As shown in table 3 of nitrite values were recorded as follows:

- 46.7% of cheese samples had values of nitrite ranging from 0 to 0.01 ppm;
- 53.3% of cheese samples had values of nitrite ranging from 0.011 to 0.1 ppm.

No evidence had values > 10 ppm that is LMA for nitrates in cheese, according to Reg. EC 1881/2006.

Statistically, nitrates had an average ( $\pm$ ) of  $0.08 \pm 0.005797$  ppm and a coefficient of variation of 56.13%. The determined values ranged between 0.0012 and 0.36 ppm.

Statistically analyzing the nitrates variation, per seasons, shows the following:

- Infall, nitrates, had an average ( $\pm$ ) of  $\pm 0.004898$  0.06721 ppm and a coefficient of variation of 28.22%.

The determined values ranged between 0.0012 and 0.078 ppm.

- In winter, nitrates, had an average ( $\pm$ ) of  $\pm 0.008054$  0.06873 ppm and a coefficient of variation of 45.39%. The determined values ranged between 0.0014 and 0.11 ppm.

- In spring, nitrates, had an average ( $\pm$ ) of  $0.1139 \pm 0.01823$  ppm and a coefficient of variation of 62.01%. The determined values ranged between 0.068 and 0.36 ppm.

- In summer, nitrates, had an average ( $\pm$ ) of  $\pm 0.005988$  0.07021 ppm and a coefficient of variation of 33.03%. The determined values ranged between 0.0012 and 0.11 ppm.

Statistically, nitrite had an average ( $\pm$ ) of  $\pm 0.002279$  0.01728 ppm and a coefficient of variation of 102.16%. The determined values ranged between 0.0 and 0,066 ppm.

Statistically analyzing the nitrite variation, per seasons, shows the following:

- In fall, nitrites, had an average ( $\pm$ ) of  $\pm 0.009867 \pm 0.003567$  ppm. The determined values ranged between 0.0 and 0.04 ppm.
- In winter, nitrites, had an average ( $\pm$ ) of  $0.01867 \pm 0.005257$ . The determined values ranged between 0.0 and 0.066 ppm.
- In spring, nitrites, had an average ( $\pm$ ) of  $\pm 0.004916 \pm 0.0268$  ppm and a coefficient of variation of 71.04%. The determined values ranged between 0.0 and 0.05 ppm.
- In summer, nitrites, had an average ( $\pm$ ) of  $\pm 0.003416 \pm 0.0138$  ppm and a coefficient of variation of 95.87%. The determined values ranged between 0.0 and 0.04 ppm. No evidence had values  $> 10$  ppm that is LMA for nitrates and nitrites in cheese.

KORÉNEKOVÁ, (2010), in one of his studies on Emmental cheeses, obtained the mean  $\text{NaNO}_2$  content in untreated and in pasteurized milk of 0.2 and 0.1  $\text{mg}\cdot\text{kg}^{-1}$ , respectively and the mean  $\text{NaNO}_3$  content of 0.9 and 0.9  $\text{mg}\cdot\text{kg}^{-1}$  respectively. He also mentioned that the milk with nitrate added had the mean content of 81.2  $\text{mg}\cdot\text{kg}^{-1}$   $\text{NaNO}_3$ . The final product had a markedly decreased content of nitrates (3.3  $\text{mg}\cdot\text{kg}^{-1}$   $\text{NaNO}_3$ ) and nitrites (0.2  $\text{mg}\cdot\text{kg}^{-1}$   $\text{NaNO}_2$ ) when compared with the values in cheese during maturation (11.3  $\text{mg}\cdot\text{kg}^{-1}$   $\text{NaNO}_3$ ; 0.4  $\text{mg}\cdot\text{kg}^{-1}$   $\text{NaNO}_2$ ). The results of his study showed that a considerable quantity of nitrates passed into the whey and that nitrates were added to the milk to prevent the blowing defect of hard cheese by micro organisms. (KORÉNEKOVÁ [13]).

## Conclusions

From the 60 investigated milk samples no evidence has exceeded the maximum permissible limit for nitrates and nitrites which is max. 10 mg/l milk. Nitrates had an average ( $\pm$ ) of  $0.0623 \pm 0.004678$  mg/l. The determined values ranged between 0.001 and 0.3 mg/l. Nitrites had an average ( $\pm$ ) of  $0.03367 \pm 0.003034$  mg/l. The determined values ranged between 0.0 and 0.08 mg/l.

Following statistical analysis of the values obtained regarding nitrate for raw milk samples and cheese final product we found statistically significant differences \*\*\* ( $p < 0.001$ ) for fall and winter samples and statistically significant \* ( $0.01 < p < 0.05$ ) for samples of spring and summer. Overall, registered statistical difference was significant \* ( $0.01 < p < 0.05$ ).

During processing nitrate concentrate into cheese 1.3 times (from 0.0623 mg/l in milk to 0.08 mg/kg in cheese).

Following statistical analysis of the values obtained regarding nitrite for raw milk samples and cheese final

product we found statistically significant differences \*\*\* ( $p < 0.001$ ) for samples taken in winter distinctively statistically significant \*\* ( $0.001 < p < 0.01$ ) for the summer samples and statistically significant \* ( $0.01 < p < 0.05$ ) for samples of autumn and spring. Overall, registered statistical difference was very significant \*\*\* ( $p < 0.0001$ ).

During processing of nitrite, content in cheese decreases by 1.9 times, from 0.03367 mg/l in milk from 0.01728 mg/kg in cheese.

## References

1. BANU, C. și VIZIREANU, CAMELIA. *Procesarea industrială a laptelui*, Editura Tehnică, București (1998).
2. BONDOC, I. *Tehnologia și controlul laptelui și produselor lactate*, vol. I, Editura „Ion Ionescu de la Brad”, Iași (2007).
3. INGA CIPROVICA, ALLA MIKELSONE. *The influence of ripening temperature on diversity of non-starter lactic acid bacteria in semi-hard cheeses*, Rom Biotechnol Lett 16 (6) Supplement (2011).
4. CRISTEA COSTACHE. *Study of the Level of Nitrates/Nitrites in Milk Products from Brașov County*, J.M.B. nr. 2, p. 28 (2011).
5. CODOBAN, JEANETA și CODOBAN, I. *Procesarea laptelui în secții de capacitate mică*, Editura Cetatea Doamnei, Piatra Neamț (2006).
6. COSTIN, G.M. și col. *Știința și ingineria fabricării brânzeturilor*, Editura Academica, Galați (2003).
7. COSTIN, G. M. *Alimente ecologice – Alimentele și sănătatea*, Editura Academica, Galați (2008).
8. GUȘ, CAMELIA. *Laptele și produsele lactate*, Ediția a II-a revizuită, Ed. Risoprint, Cluj-Napoca (2005).
9. GUZUN, V. și col. *Industrializarea laptelui*, Editura Tehnică-Info, Chișinău (2001).
10. JIMBOREAN, MIRELA, ȚIBULCĂ, D. *Tehnologia de fabricare a brânzeturilor*, Editura Risoprint, Cluj-Napoca (2006).
11. MIRELA JIMBOREAN, D. ȚIBULCĂ, ADRIANA PĂUCEAN. *Researches of Nitrates Content in Raw Milk*, Buletin USA-MV Cluj-Napoca, vol. 68 (2), p. 540 (2011).
12. MIRELA JIMBOREAN, D. ȚIBULCĂ. *Seasonal Variation of Nitrates And Nitrites In Milk As Raw Material*, *Analele Universității din Craiova*, vol. XVII(LIII) (2012).
13. KORÉNEKOVÁ B, KOTTFFEROVÁ J, KORÉNEK M. *The fate of added nitrate used in the manufacture of Emmental cheese*, Food Addit Contam., 17(5):373-7 (2010).
14. CRINA MUREȘAN, MARIA TOFANĂ, SONIA SOCACI, LIANA SALANȚĂ. *Content Evaluation on*

- Nitrates and Nitrites in Local and Imported Vegetables*, Bulletin UASVM Agriculture, 69(2), p. 510 (2012).
15. SERAPHIM KR<sup>1</sup>, de SIQUEIRA ME. *Nitrates and Nitrites in Homemade and Industrial Cheeses commercialized in the Southern region of Minas Gerais, Brazil*, Arch Latinoam Nutr.; 50(1):87-90 (2000).
  16. VASILICA SIMION, Gh. CÂMPEANU, GINA VASILE, MIHAELA ARTIMON, LUMINIȚA CATANĂ, MIOARA NEGOIȚĂ. *Nitrate and nitrite accumulation in tomatoes and derived products*, Rom Biotechnol Lett, 13 (4): 3785-3790 (2008).
  17. SUNGUR Ş, ATAN MM. *Determination of Nitrate, Nitrite and Perchlorate Anions in Meat, Milk and Their Products Consumed in Hatay Region in Turkey*, Food Addit Contam Part B Surveill, 6(1):6-10 (2013).
  18. ŞINDILAR, E. *Controlul igienic al produselor și subproduselor de origine animală*, vol. 1, 2, Editura I.N.R.C.S., Iași (2000).
  19. ALI TOPÇU, AYLIN AYAZ TOPÇU, ILBILGE SALDAMLI and MINE YURTTAGÜL. *Determination of nitrate and nitrite content of Turkish cheeses*, African Journal of Biotechnology, vol. 5 (15), pp. 1411-1414 (2006).
  20. L. TUDOR, ELENA MITTRĂNESCU, LAURA TUDOR, F. FURNARIS. *Assessment of Nitrate and Nitrite Content of Romanian Traditional Cheese*, Lucrări Științifice Medicină Veterinară, vol. XL, Timișoara (2007).
  21. ȚIBULCĂ AURORA. *Cercetări privind influența factorilor tehnologici și de igienă asupra calității cașcavalurilor fabricate în zona de nord a Transilvaniei*, Cluj-Napoca (2011).
  22. ȚIBULCĂ, D. și MIRELA A. JIMBOREAN. *Fabricarea produselor lactate și a brânzeturilor*, Editura AcademicPres, Cluj-Napoca (2003).
  23. WALKER R. *Nitrates, Nitrites and N-nitrosocompounds: a Review of the Occurrence in Food and Diet and the Toxicological Implications*, Food Addit Contam. Nov-Dec; 7(6): 717-68 (1990).