



Received for publication, July, 15, 2017
Accepted, August, 20, 2018

Original paper

Seed germination of some medicinal plant species for conservative purpose

RODICA CATANĂ^{1*}, MĂDĂLINA LAZĂR², IRINA HOLOBIUC¹,
LARISA FLORESCU¹

¹Institute of Biology Bucharest, Romanian Academy, 296 Splaiul Independenței St., 060031, P.O. Box 56-53, Bucharest, Romania

²University of Bucharest, Faculty of Biology, 1-3 Portocalelor Entrance, sector 6, Bucharest, România

Abstract

Due to the overexploitation, many medicinal plants are in decline. Enhancing the seed germination and the vigorous seedlings production is crucial for their cultivation and for ex situ conservation. Seven medicinal plant species were studied concerning seed germination as effect of different treatments (sterilization procedures and stimulation of germination). Seed sterilization with HgCl₂ 0.1% for 12 minutes was effective for controlling microbial and fungal contamination in all tested medicinal species. *In vitro* conditions were efficient for seed germination for 4 species. The stimulation treatments tested (KNO₃ 0.1%, NaCl 0.5% and GA₃ 0.5%) determined an improvement of seed germination capacity compared with untreated seeds.

Keywords

Medicinal plant, germination, stimulation, conservation.

To cite this article: CATANĂ R, LAZĂR M, HOLOBIUC I, FLORESCU L.
Seed germination of some medicinal plant species for conservative purpose. *Rom Biotechnol Lett.* 2020; 25(3): 1621-1627. DOI: 10.25083/rbl/25.3/1621.1627

✉ *Corresponding author: RODICA CATANĂ, Institute of Biology Bucharest, Romanian Academy, 296 Splaiul Independenței St., 060031, P.O. Box 56-53, Bucharest, Romania
E-mail: catanarodica@yahoo.com

Introduction

The medicinal plant species represent a consistent part of the natural biodiversity. In May 2015, an IUCN report showed that Europe's medicinal plants are in decline due to collection of plants from the wild (for their medicinal value, ornamental and horticultural plant trade) (ALLEN & al [1]). In our days, ~ 80% of the people use medicinal plant species (crude materials, processed products like pharmaceuticals, herbal remedies, teas, spirits, cosmetics, sweets, dietary supplements, insecticides etc.) for their primary health care (ARCEUSZ & al [2]).

In România almost 900 plant species represent medicinal flora (MUNTEAN [3]). The research directions in these species were focused on their biology, the content of active ingredients and the improvement of culture technology (MUNTEAN [3]).

Three from the selected species are included in the European Red List of Medicinal Plants like least concern (*Agrimonia eupatoria*, *Cnicus benedictus*, *Geum urbanum*) (ALLAN et al [1]). The others, *Foeniculum vulgare* and *Hyssopus officinalis* are plants produced at commercial scale. *F. vulgare* is characterised by poor seed germination (TAHA EI et al [4]).

According to Guidelines on the conservation of medicinal plants [5], the strategies used for the conservation of medicinal plant species applied over the world are represented by *in situ* and *ex situ* strategies. The *ex situ* conservation was classified as *in vivo* (botanical gardens, field cultures, seed banks) and *in vitro* (tissue cultures, artificial seeds, cryopreservation) methods. The plant cells and tissues culture technologies offer a great opportunity to exploit the medicinal plants (SHARMA & al [6]).

Seed germination and seedling establishment are the initial stages of plant development. These processes are unpredictable over space and time. The germination process is specific for each species, depending on different factors like seed maturity, physical conditions (temperature and humidity), harvesting methods, transport and maintenance. In the case of medicinal plants, there is the possibility that seeds can germinate only in the native environment and fail to germinate under the laboratory conditions (GUPTA [7]). Knowledge of the seeds germination of medicinal plants has proved to be useful in development of appropriate conservation strategies (KANDARI & al [8]).

Low germination rates (frequently result from fungal infection or mechanical damage) or specific ecological requirements may affect the cultivation of some herbs

(VINES [9]). Enhancing the seeds germination and developing of the vigorous seedlings is crucial for an *ex situ* conservation strategy. The first step of *in vitro* cultures establishment is the sterilization – process consisting in different treatments having as effect explants free of contamination; viable decontaminated explants are used further for the initiation of tissues culture. When in case of conventional methods, a low rate or no germination was registered, *in vitro* techniques may enhance the germination process (FAY [10]).

The aim of the present study was to check and improve the seed germination capacity of some medicinal plant for conservative purpose.

Materials and Methods

Mature seeds from different species of medicinal plants belonging to four families (Table 1) were tested concerning the germination capacity.

The seeds of *F. vulgare*, *P. anisum* and *H. officinalis* were bought from market, in 2014 and others (*A. eupatoria*, *C. benedictus*, *G. urbanum*, *C. sativum*) were kindly provided by Nicoleta Constantin in 2012, from Botanical Garden, Bucharest.

All seeds were hand sorted to select undamaged seeds. For germination experiments were used 75 seeds in 3 replications/ each treatment. After treatments, seeds were inoculated in distilled water, on filter paper in Petri dishes in culture chamber at 25°C and a photoperiod of 16/8 and illumination at 1500 lux. Germinated seeds were counted every day. A seed was considered germinated when the tip of the radicle had grown free of the seed coat.

Experiment 1: Seed germination *in vitro* and *in vivo*

To compare the seed germination capacity in 7 medicinal plant species, *in vitro* and *in vivo* conditions were tested, temperature and illumination regime being the same (25°C and 16/8 hours photoperiod, 1500 lux illumination).

For *in vitro* germination, all seeds were sterilized. Steps of sterilization protocol consisted in washing in running tap water (for 2-3 hours), the immersion in ethanol 70° (30 seconds), in the treatment with HgCl₂ 0.1%, (for 12 minutes) followed by three washing in sterile distilled water.

For *in vivo* germination, unsterilized seeds were placed on filter paper discs wetted with distilled water in placed in 10 cm Petri dishes.

Table 1. The medicinal plant species tested for seed germination capacity

Plant species	Family	Life cycle	Used for	Seeds aspect
<i>Agrimonia eupatoria</i> L.	Rosaceae	perennial	disorders related to the liver, bile, gastrointestinal, respiratory tract	
<i>Cnicus benedictus</i> L.	Asteraceae	annual	<i>In vitro</i> antibacterial activity against <i>Bacillus subtilis</i> , <i>Brucella species</i> , <i>E. coli</i> , <i>Proteus species</i> , <i>P. aeruginosa</i> , <i>S. aureus</i> , <i>Streptococcus faecalis</i> , anti-HIV, anticancer agents	
<i>Geum urbanum</i> L.	Asteraceae	perennial	rheumatism, gout, infections, fever	
<i>Coriandrum sativum</i> L.	Apiaceae	annual	cold, seasonal fever, stomach disorders, against worms, rheumatism and pain in the joints	
<i>Foeniculum vulgare</i> Mill.	Apiaceae	perennial	antioxidant, anti-inflammatory, antimicrobial, estrogenic, diuretic, antithrombotic, hepatoprotective, memory enhancing, antimutagenic	
<i>Pimpinella anisum</i> L.	Apiaceae	annual	analgesic, carminative, aromatic, disinfectant, diuretic	
<i>Hyssopus officinalis</i> L.	Lamiaceae	perennial	expectorant, antiseptic	

Experiment 2: Seed germination improvement

Among the species analyzed, were chosen two species, one with no germination *in vivo* and low *in vitro* response (*C. sativum*) and one without any positive germination response (*C. benedictus*). For these species, some stimulation treatments were applied to improve the seeds germination. Seeds were maintained in KNO₃ 0.1%, NaCl 0.5% and GA₃ 0.5% solutions for 3 days at room temperature. The control was represented by seeds cultured on filter paper with distilled water for 3 days without any treatment. No sterilization protocol was used.

After stimulation of germination during 3 days, the seeds (treated and untreated) were inoculated in a substrate

consisted in ground and perlite mix in equal parts. Seeds were periodically wetted with tap water without any growth factors or stimulants.

The recorded parameters were the final germination percentage (mean value \pm standard error) and mean germination time calculated after formulas:

Final germination percentage = total number of germinated seeds/total number of seeds X 100.

Mean germination time = $\Sigma (nT) / \Sigma N$ (ELLIS & ROBERTS [11]), where n = number of seeds newly germinated at time T at 25°C; T = time (hours, days) from the beginning of the germination test, N = final germination.

The plants obtained from germinated seeds were further used as sources of explants for *in vitro* multiplication.

The photos were taken with a Panasonic DMC-LS80 camera. The results were statistically analyzed using ANOVA from Daniel's XL Toolbox version 6.52 (<http://xltoolbox.sourceforge.net>). A posthoc test (Bonferonni-Holm) was applied to check the significant differences among means.

Results and Discussion

In the present study, aspects concerning seeds germination in seven medicinal plants were evaluated. In the first experiment, seeds were sterilized. It is known that seeds collected from the natural habitats are often contaminated with exogenous and endogenous microbial contaminants (HALLOIN [12]; AHMAD & al [13]). The seeds sterilization is a prerequisite for successful *in vitro* germination. Agents as ethanol, sodium hypochlorite,

calcium hypochlorite, chlorine gas and mercuric chloride have been widely used for surface sterilization of parts of the plant and seed material of various species (TALEI & al [14]; DAUD & al [15]; BARAMPURAM & al [16]).

In our experiment, although HgCl₂ is a toxic agent, at the level used (0.1%) and time of exposure (12 minutes) was effective for controlling microbial and fungal contamination in all medicinal plant species tested. Previous works reported that treatment with HgCl₂ at 0.1-0.2% for 10-15 minutes was effective for surface sterilization of other like *Aquilaria agallocha* (HE & al [17]) and *Aquilaria sinensis* (SHU & al [18]).

Germination percentage recorded *in vivo* and *in vitro* conditions (Table 2) shows that *in vitro* culture conditions had a stimulatory effect in the case of *Agrimonia eupatoria*, *Geum urbanum*, *Coriandrum sativum* and *Pimpinella anisum* seeds.

Table 2. Germination percentage of the tested medicinal plant species *in vitro* and *in vivo* conditions

Medicinal plant species	Treatments	Germination percentage	F	P
<i>Agrimonia eupatoria</i>	<i>in vivo</i>	0 ^a	57.558	3.20549E-12
	<i>in vitro</i>	40.9 ^b		
<i>Geum urbanum</i>	<i>in vivo</i>	0 ^a	57.558	3.20549E-12
	<i>in vitro</i>	43.47 ^b		
<i>Hyssopus officinalis</i>	<i>in vivo</i>	16.66 ^a	14.062	0.000
	<i>in vitro</i>	0 ^b		
<i>Coriandrum sativum</i>	<i>In vivo</i>	0 ^a	11.363	0.000
	<i>in vitro</i>	13.63 ^b		
<i>Foeniculum vulgare</i>	<i>In vivo</i>	44 ^a	28.971	2.77284E-07
	<i>in vitro</i>	7.14 ^b		
<i>Pimpinella anisum</i>	<i>In vivo</i>	0 ^a	6,428	0,012254
	<i>in vitro</i>	8.82 ^b		
<i>Cnicus benedictus</i>	<i>In vivo</i>	0		
	<i>in vitro</i>	0		

Values having the same letter are not significantly different at 0.05 significance level.

The final germination percentage varied between 7.14% for *F. vulgare* and 43.4% for *G. urbanum*. Using Bonferonni-Holm test showed significant differences between sterilized and unsterilized seeds in the case of *Hyssopus officinalis*, *Coriandrum sativum*, *Foeniculum vulgare* (p<0.0002), *Pimpinella anisum* (at p<0.01) (Table 2). Seeds of *Cnicus benedictus* did not germinated in the first experiment.

This lower germination percentage could be attributed to factors such as lack of nutrients, seeds being inoculated in distilled water without growth factors or germination enhancing factors.

Seeds inoculated *in vivo*, at the same temperature, humidity, illumination and substrate, but without sterilization, did not germinate in five tested species. Only in the case of *F. vulgare* (44%) and *H. officinalis* (16.66%) the seeds germinated.

Our results concern the efficiency of *in vitro* conditions than *in vivo* are in accordance with studies about *Cassia alata* seed germination (THIRUPATHI & al [19]). *Asparagus racemosus* (RAGHAV & KASERA [20]) and *Alstroemeria ligtu* hybrid (NASRI & al [21]) seeds germination was improved by *in vitro* conditions.

In the case of fennel seeds, it was demonstrated that the optimum temperature for germination is around 22°C with small differences for each variety (THOMAS [22]). It was previously reported that the fennel seeds need some treatments to increase the germination rate (NEACŞU & al [23]). In our experiment, seed germination was low both *in vivo* (44%) and *in vitro* (7.14%) (Table 2). In the case of this species, is possible that sterilization with HgCl₂ inhibited the seed germination.

Usually, *Agrimonia eupatoria* seeds germinate in 2-6 weeks at 13°C (RICE [24]). In our experiment, seeds

were collected in 2012 and were stored at room temperature for 2 years. No seed germination was observed at 25°C, neither control seeds nor seeds treated with stimulatory treatments. After *in vitro* procedures (sterilization with ethanol and HgCl₂ at 25°C) 40.9% of *Agrimonia* seeds germinated. An explanation of *in vitro* seed germination in this species might be that sterilization with ethanol and HgCl₂ break the dormancy of zygotic embryos. These results are accordingly with others authors (JEAVONS

& JARVIS [25]) which used a treatment with ethanol and mercuric chloride to break the dormancy of intact hazel seeds.

In the second experiment, the effects of stimulatory treatments (KNO₃, NaCl and GA₃) applied *in vivo* were analyzed (Table 3). The stimulation treatments have generally been applied to enhance seed germination (JOSHI & DHAR [26]) and to increase seedling vigor (LEITE & al [27]).

Table 3. Seeds germination and applied treatments for *C. benedictus* and *C. sativum*

Medicinal plant species	Treatments	Mean germination
<i>Cnicus benedictus</i>	Control	0±0 ^a
	KNO ₃ 0.1%	0.671±0.47 ^b
	NaCl 0.5%	0.315±0.46 ^c
	GA ₃ 0.5%	0.644±0.48 ^c
<i>Coriandrum sativum</i>	Control	0±0 ^a
	KNO ₃ 0.1%	0.671±0.47 ^b
	NaCl 0.5%	0.63±0.44 ^c
	GA ₃ 0.5%	0.329±0.47 ^c

Values having the same letter are not significantly different at 0.05 significance level.

In our experiments, the treatments with KNO₃, NaCl or GA₃ were effective in improving seed germination. Seeds of *Coriandrum sativum* exposed at KNO₃ 0.1% and GA₃ 0.5% showed an improved germination capacity (68%) comparing with control. In the case of *Cnicus benedictus* the highest final germination percentage (67%)

was recorded for seeds soaked in 0.5% GA₃. A lower germination (33.33%) was recorded in the case of seeds treated with 0.5% NaCl. The results showed that there are significant differences between treated and untreated seeds for both species *Coriandrum sativum* and *Cnicus benedictus* (Table 4).

Table 4. Result of posthoc test, showing the significance differences between stimulation treatments (using Daniel's XL Toolbox version 6.52).

		<i>Coriandrum sativum</i>		<i>Cnicus benedictus</i>	
		Critical	P	Critical	P
Control	KNO ₃ 0.1%	0.008	0.000	0.008	0.000
Control	GA ₃ 0.5%	0.01	0.000	0.01	0.000
Control	NaCl 0.5%	0.016	0.000	0.0125	0.000
KNO ₃ 0.1%	NaCl 0.5%	0.012	0.000	0.016	0.000
NaCl 0.5%	GA ₃ 0.5%	0.05	0.377	0.025	0.000
KNO ₃ 0.1%	GA ₃ 0.5%	0.025	0.000	0.05	0.734

There are several studies concerning seed germination improved with KNO₃ and NaCl solutions in some medicinal plants (GUPTA [7]; SHIVKUMAR & al [28]; KAMBIZIA & al [29]; KULKARNI & al [30]; FARIMAN & al [31]).

The shortest mean germination time was registered in the case of *Cnicus benedictus* treated with 0.5% GA₃ (13.5 days). With the same treatment, seeds of *Coriandrum sativum* starts to germinated after 16 days. After

treatment with 0.5% NaCl, germination capacity starts after 36 days.

Our results are in accordance to other studies which showed that potassium nitrate treatment increased seeds germination in *Avena fatua* (HILTON [32]), *Hypericum aviculariifolium* (ÇIRAK & al [33]), chicory (TZORTZAKIS [34]), *Foeniculum vulgare* and *Cuscuta epithimum* (TAVILI & al [35]).

Conclusion

Our results showed that controlled conditions assured by *in vitro* techniques were efficiently for seed germination for almost species tested. Seeds of *Foeniculum vulgare* may be sensible at HgCl₂, *in vitro* germination being inhibited. *Cnicus benedictus* seeds did not germinate *in vivo* and *in vitro* conditions, only stimulatory treatments determined an improvement of the germination capacity. Treatment with KNO₃ positively influenced the germination of *Coriandrum sativum* and *Cnicus benedictus*.

Acknowledgements

The study was funded by project no. RO1567-IBB06/2015 from the Institute of Biology Bucharest of Romanian Academy and is part of the Master degree thesis of Miss Lazăr Mădălina.

References

1. D. ALLEN, M. BILZ, D.J. LEAMAN, R.M. MILLER, A. TIMOSHYNA, J. WINDOW. European Red List of Medicinal Plants. Luxembourg: Publications Office of the European Union (2014).
2. A. ARCEUSZ, I. RADECKA, M. WESOŁOWSKI. Identification of diversity in elements content in medicinal plants belonging to different plant families. *Food Chemistry*, 120: 52-58 (2010).
3. L.S. MUNTEAN. One hundred ten years of research on medicinal plants, at the agronomy in Cluj (1904-2014). *Hop and Medicinal Plants*, XXII (1-2): 7-16 (2014).
4. A. TAHAEI, A. SOLEYMANI, M. SHAMS. Seed germination of medicinal plant, fennel (*Foeniculum vulgare* Mill), as affected by different priming techniques. *Applied Biochemistry and Biotechnology*, 180(1): 2-40 (2016).
5. Guidelines on the Conservation of Medicinal Plants, Published by The International Union for Conservation of Nature and Natural Resources (IUCN), Gland, Switzerland, in partnership with The World Health Organization (WHO), Geneva, Switzerland, and WWF – World Wide Fund for Nature, Gland, Switzerland (1993).
6. S. SHARMA, N. RATHI, B. KAMAL, D. PUNDIR, B. KAUR, S. ARYA. Conservation of biodiversity of highly important medicinal plants of India through tissue culture technology – a review. *Agriculture and biology journal of America*, 1(5): 827-833 (2010).
7. V. GUPTA. Seed germination and dormancy breaking techniques for indigenous medicinal and aromatic plants. *Journal of Medicinal and Aromatic Plants Science*, 25: 402-407 (2003).
8. L.S. KANDARI, K.S. RAO, K. CHAUHAN, R.K. MAIKHURI, V.K. PUROHIT, P.C. PHONDANI, K.G. SAXENA. Effect of presowing treatments on the seed germination of two endangered medicinal herbs of the Himalaya (*Angelica glauca* Edgew and *Pleurospermum angelicoides* Wall. Ex DC. Benth. Ex C.B. Clarke). *Proc. Indian Nat. Sci. Acad.*, 73: 11-16 (2007).
9. G. VINES. Herbal harvests with a future: towards sustainable sources for medicinal plants. *Plantlife International* (2004).
10. M.F. FAY. Conservation of rare and endangered plants using *in vitro* methods. *In vitro Cell. Dev. Biol.*, 28: 1-4 (1992).
11. R.H. ELLIS, E.H. ROBERTS. Towards a rational basis for testing seed quality. In *Seed Production*, Hebblethwaite P.D. eds., Butterworths, London, (1980).
12. J.M. HALLOIN. Postharvest infection of cottonseed by *Rhizopus arrhizus*, *Aspergillus niger*, *Aspergillus flavus*. *Phytopathology*, 65: 1229-1232 (1975).
13. F. AHMAD, G.J. DAGLISH, A.W. RIDLEY, G.H. WALTER. Responses of *Tribolium castaneum* to olfactory cues from cotton seeds, the fungi associated with cotton seeds, and cereals. *Entomol. Exp. Appl.*, 145: 272-281 (2012).
14. D. TALEI, A. VALDIANI, M.P. ABDULLAH, S.A. HASSAN. A rapid and effective method for dormancy breakage and germination of King of Bitters (*Andrographis paniculata* Nees.) seeds. *Maydica*, 57(2): 98-105 (2012).
15. N.H. DAUD, S. JAYARAMAN, R. MOHAMED. An improved surface sterilization technique for introducing leaf, nodal and seed explants of *Aquilaria malaccensis* from field sources into tissue culture. *Asian Pac. J. Mol. Biol. Biotechnol.*, 20: 55- 58 (2012).
16. S. BARAMPURAM, G. ALLEN, S. KRASNYSKI. Effect of various sterilization procedures on the *in vitro* germination of cotton seeds. *Plant Cell. Tiss. Organ. Cult.*, 18: 179-185 (2014).
17. M.L. HE, S.Y. QI, L.J. HU. Rapid *in vitro* propagation of medicinally important *Aquilaria agallocha*. *Journal of Zhejiang University Science*, 6(8): 849-852 (2005).
18. Y.Q. SHU, L.H. MENG, D.L. LI, H.Z. CHUAN, J.H. LAN, Z.Z. HUI. Production of 2-(2-phenylethyl) chromones in cell suspension cultures of *Aquilaria sinensis*. *Plant Cell. Tiss. Org.*, 83: 217-221 (2005).
19. M. THIRUPATHI, M.K. KUMAR, D. SRINIVAS, K. RAJENDAR, K.R. JAGANMOHAN. A study on seed germination of *Cassia alata* Linn an anti-allergenic plant. *Journal of Phytochemistry*, 4(5): 13-15 (2012).
20. A. RAGHAV, P.K. KASERA. Seed germination behavior of *Asparagus racemosus* (Shatavari) under

- in-vivo* and *in-vitro* conditions. *Asian Journal of Plant Science and Research*, 2(4): 409-413 (2012).
21. F. NASRI, N. GHADERI, J. MOHAMMADI, S.N. MORTAZAVI, K.S. MAHMOOD. The effect of gibberellic acid and stratification on germination of *Alstroemeria* (*Alstroemeria ligtu* hybrid) seed under *in vitro* and *in vivo* conditions. *Journal of Ornamental Plants*, 3(4): 221-228 (2013).
 22. T.H. THOMAS. Responses of Florence fennel (*Foeniculum vulgare azoricum*) seeds to light, temperature and gibberellin A_{4/7}. *Plant Growth Regul.*, 14: 139-143 (1994).
 23. V.I. NEACȘU, A.S. APAHIDEAN, A.M. HUSTI, R. CICEVAN. Study of fennel seed germination (*Foeniculum vulgare* var. *Azoricum*). *Agriculture – Science and Practice*, 1(2): 19-22 (2015).
 24. G. RICE. Editor, Growing from Seed. Volume 1. Publisher: Thompson and Morgan (1987).
 25. R.A. JEAVONS, B.C. JARVIS. The breaking of dormancy in hazel seed by pretreatment with ethanol and mercuric chloride. *New Phytol.*, 96: 551-554 (1984).
 26. M. JOSHI, U. DHAR. Effect of various pre-sowing treatments on seed germination of *Heracleum candicans* Wall. Ex DC: a high value medicinal plant. *Seed Sci. Technol.*, 31: 737-743 (2003).
 27. V.M. LEITE, C.A. ROSOLEM, J.D. RODRIGUES. Gibberellin and cytokinin effects on soybean growth. *Sci. Agric.*, 60: 537-541 (2003).
 28. V. SHIVKUMAR, R. ANANDLAKSHMI, R.R. WARRIER, M. TIGABU, P.C. ODEN, S.N. VIJAYACHANDRAN, S. GEETHA, B.G. SINGH. Effect of presowing treatments, desiccation and storage conditions on germination of *Strychnos nux-vomica* seeds, a valuable medicinal plant. *New Forest*, 32: 121-131 (2006).
 29. L. KAMBIZIA, P.O. ADEBOLAB, A.J. AFOLAYANA. Effects of temperature, pre-chilling and light on seed germination of *Withania somnifera*, a high value medicinal plant. *S. Afr. J. Bot.*, 72: 11-14 (2006).
 30. M.G. KULKARNI, R.A. STREET, J. VAN STADEN. Germination and seedling growth requirements for propagation of *Dioscorea dregeana* (Kunth) Dur. and Sching – a tuberous medicinal plant. *S. Afr. J. Bot.*, 73: 131-137 (2007).
 31. Z.K. FARIMAN, M. AZIZI, S. NOORI. Seed germination and dormancy breaking techniques for *Echinacea purpurea* L. *J. Biol. Environ. Sci.*, 5(13): 7-107 (2011).
 32. J.R. HILTON. The influence of light and potassium nitrate on the dormancy and germination of *Avena fatua* L. (wild oat) seed and its ecological significance. *New Phytologist*, 96(1): 31-34 (1984).
 33. C. ÇIRAK, K. KEVSEROĞLU, A.K. AYAN. Breaking of seed dormancy in a Turkish endemic *Hypericum* species: *Hypericum aviculariifolium* subsp. *Depilatum* var. *depilatum* by light and some pre-soaking treatments. *J. Arid. Environ.*, 68(1): 159-164 (2007).
 34. N.G. TZORTZAKIS. Effect of pre-sowing treatment on seed germination and seedling vigor in endive and chicory. *Hort. Sci.*, 36(3): 117-125 (2009).
 35. A. TAVILI, A. FARAJOLLAHI, H. POUZESH, E. BANDAK. Treatment induced germination improvement in medicinal species of *Foeniculum vulgare* miller and *Cuscuta epithimum* (L.) L. *Journal of Modern Applied Science*, 4(7): 163-169 (2010).